

SYNTHESIS, CHARACTERIZATION AND ANTIMICROBIAL ACTIVITY STUDIES OF 5,5'-(6-(METHYL(PHENYL)AMINO)-1,3,5-TRIAZINE-2,4-DIYL) BIS(AZANEDIYL)DIQUINOLIN-8-OL AND THEIR CO-ORDINATION POLYMERS

Ankita I.Chaudhari¹ and JA.Chaudhari²

¹Department of Chemistry, Research Scholar of JTT University, Rajasthan-333 001, India.

²Shri R.K.Parikh Arts & Science college Petlad,Gujarat-388450, India.

ABSTRACT


Co-ordination polymers containing a novel bis ligand namely 5,5'-(6-(methyl(phenyl)amino)-1,3,5-triazine-2,4-diyl)bis(azanediy)diquinolin-8-ol(TBQ-2) have been prepared with metal ion like Zn⁺², Cu⁺², Ni⁺², Co⁺² and Mn⁺². The novel bis bidentate ligand (TBQ-2) is synthesized by condensation of 5-amino-8-hydroxyquinoline with 4,6-dichloro-N-methyl-N-phenyl-1,3,5-triazin-2-amine in presence of base catalyst. All of these co-ordination polymers and parent ligand were characterized by elemental analysis, IR spectra and diffuse reflectance spectral studies for their structure determination. The thermal stability was evaluated by thermo gravimetric analyses (TGA). In addition, all of the coordination polymers have been characterized by their magnetic susceptibilities. The microbicidal activity of all the samples have been monitored against plant pathogens.

Keywords: 8-hydroxyquinoline, antibacterial and antifungal activities, coordination polymers.

INTRODUCTION

Nitrogen containing heterocyclic play an important role, not only for life science industries but also in many other industrial fields related to special and fine chemistry. Among them s-triazine ring containing derivatives have been reported for applicable mostly as reactive dyes and some are used as polymers and drugs .The study of co-ordination polymers has made much progress¹⁻³. 5-amino 8-quinolinol is well known as an analytical reagent^{4,5}. Its various derivatives are very useful in pharmaceuticals⁶. Several azo dyes based on 8-quinolinol are also reported for dyeing of textiles as well as their chelating properties^{7,8}. A promising method has been reported for the formation of coordination polymers of enhanced chelating ability by using a bidentate

8-hydroxyquinoline moiety in which two 8-hydroxyquinolinyl end groups are joined with bridge, usually at the 5,5'-position⁹⁻¹¹. The 5-Amino 8-hydroxy quinolinol is the easiest preparable precursor for the preparation of bis-ligand and thus bis-ligands based on 5-Amino 8-hydroxyquinoline have been reported for coordination polymers^{11,12}.ion exchange resins have also been prepared from 5-amino 8-hydroxyquinoline and amino or hydroxyl functionalized polymers¹³⁻¹⁶. We are also synthesized compound of TBQ-2. Hence, In this paper, we report newly compound of in continuous of this work¹³ the present paper deals with synthesis, characterization and chelating properties of ligand (TBQ-2) and its co-ordination polymers are shown in Scheme 1.



EXPERIMENTAL

MATERIALS

All the chemicals used were of pure grade. 5-amino 8-hydroxyl quinoline was obtained from local dealer.

Synthesis of 5,5'-(6-(methyl(phenyl)amino)-1,3,5-triazine-2,4-diyl)bis(azanediyl)diquinolin-8-ol (TBQ-2)

To a suspension of 5-amino 8-hydroxyquinoline (3.2g, 0.02 mol), 4,6-dichloro-N-methyl-N-phenyl-1,3,5-triazin-2-amine (2.54 g, 0.01 mol) in an acetone-water mixture was added. Then K_2CO_3 (0.02 mol) was added as an acid accepted¹⁷. The resulting mixture was refluxed for 3 hr with occasional shaking. The resulting suspension, which contained a precipitate, was neutralise and then filtered. The solid product was collected and dried to give TBQ-2 (66% yield). The product melted with decomposition at above 255°C (uncorrected).

Synthesis of coordination polymer

A solution of metal (0.01 mol) in aqueous formic acid was added drop wise to a solution of TBQ-2 (0.01 mol) in aqueous formic acid with stirring. The reaction mixture was heated on a water bath for 1.0hr. The reaction mixture was made alkaline by the addition of dilute aqueous ammonia until the precipitation was completed. The polymer separated out in the form of a suspension and was digested on a boiling water

bath for about 1 hr. Finally, the resultant solid was collected by filtration and washed with hot water, dimethylformamide (DMF), and then acetone. The polymer [TBQ-2 -M⁺²] (resultant product) was air-dried.

Antimicrobial Activities

Antibacterial activity and antifungal activities of TBQ-2 ligand and its coordination polymers were studied against gram-positive bacteria (*Bacillus subtilis* and *staphylococcus aureus*) and gram-negative bacteria (*E.coli*, *salmonella typhi* and *klebsiella promioe*) and plant pathogenic organisms used were *Aspergillus niger*, *Candida albicans*, *Trichoderma harsianum*, *Mucor mucedo*, and *Botrytis cinerea* at a concentration of 50 µg/ml by agar cup 520 plate method. The methanol system was used as control in this method. The area of inhibition of zone was measured in mm.

MEASUREMENT

The C, H, N contents of metal were determined by TF-Flash-1101 EA. The metals contents of metal chelates were determined volumetrically by Vogel's method¹⁸. To a 100mg chelate sample, 1ml of HCl, H₂SO₄ and HClO₄ each were added and then 1 gm of NaClO₄ was added. The mixture was evaporated to dryness and the resulting salt was dissolved in double distilled water and diluted to the mark. From this solution the metal content was determined by

titration with standard EDTA solution. Infrared spectra of the synthesized compounds were recorded on Nicolet 760 FT-IR spectrometers. NMR spectrum of ligand was recorded on a Bruker spectrophotometer at 400 MHz. Magnetic susceptibility measurement of the synthesized coordination polymer was carried out on Gouy Balance at room temperature. The electronic spectra of coordination polymer in solid were recorded at room temperature. MgO was used as a reference. Antimicrobial activity of all the samples was monitored against various gram positive (+) and gram negative(-) organisms, following the method reported in the literature [19,20].

RESULTS AND DISCUSSION

The synthesis of 5,5'-(6-(methyl(phenyl)amino)-1,3,5-triazine-2,4-

diyl)bis(azanediy) diquinolin-8-ol (TBO-2) was performed by a simple nucleophilic substitution reaction of 4,6-dichloro-N-methyl-N-phenyl-1,3,5-triazin-2-amine and 5-amino 8-hydroxy quinoline. The resulting TBO-2 ligand was an amorphous color powder. The ligand is characterized by elemental analysis as well as ¹H NMR and IR spectroscopic techniques as given below.

Elemental Analysis

The analytical and physical properties of the ligand and its coordination polymers are listed in Table-1. and are also consistent with the predicted structure. The results show that the metal: ligand (M: L) ratio for all divalent metal chelate is 1:1.

Table 1: Analysis of TBO-2 ligand and their co-ordination polymers

Empirical Formula	Mol. Cal g/mol	Yield %	Elemental Analysis (%) Found(Calcd)			
			C	H	N	M
C ₂₈ H ₂₂ N ₈ O ₂	502	66	66.93 (66.90)	4.30(4.38)	22.30(22.31)	--
C ₂₈ H ₂₀ N ₈ O ₂ Cu.2H ₂ O	599	75	56.00 (56.04)	4.00(4.00)	18.68(18.60)	10.55(10.59)
C ₂₈ H ₂₀ N ₈ O ₂ Ni.2H ₂ O	591	70	56.80 (56.85)	4.00(4.06)	19.00(18.95)	09.30(09.30)
C ₂₈ H ₂₀ N ₈ O ₂ Co.2H ₂ O	595	77	56.40 (56.47)	4.00((4.03)	18.80(18.82)	09.90(09.91)
C ₂₈ H ₂₀ N ₈ O ₂ Mn.2H ₂ O	595	75	56.45 (56.47)	4.00(4.03)	18.85(18.82)	09.90(09.91)
C ₂₈ H ₂₀ N ₈ O ₂ Zn.2H ₂ O	601	75	55.90 (55.90)	4.00(4.00)	18.60(18.63)	10.80(10.81)

Table 2: spectral features and magnetic moment of co-ordination polymers

Metal Chelates	BM	Electronic Spectral Data cm ⁻¹	Transitions	IR spectral features Common for all cm ⁻¹
TBO-2 -Cu ⁺²	2.05	22500 15590	C.T ² E _g → ² T _{2g}	3300 Quinoline Moiety 2300 1610 1470 1100 C-O-M & 1280 O-M 710 M-N 550
TBO-2 -Ni ⁺²	2.82	22580 15100 8300	³ A _{2g} → ³ T _{1g} (P) ³ A _{2g} → ³ T _{1g} (F) ³ A _{2g} → ³ T _{2g}	
TBO-2 -Co ⁺²	4.87	21270 18600 11950	⁴ T _{1g} (F) → ⁴ A _{2g} ⁴ T _{1g} (F) → ⁴ T _{1g} (P) ⁴ T _{1g} (F) → ⁴ T _{2g}	
TBO-2 -Mn ⁺²	5.91	22450 17620 14800	⁶ A _{1g} → ⁶ A _{1g} (⁴ E _g) ⁶ A _{1g} → ⁴ T _{2g} (⁴ G) ⁶ A _{1g} → ⁴ T _{1g} (⁴ G)	
TBO-2 -Zn ⁺²	Diamagnetic		-----	

IR Analysis

The important infrared spectral bands and their tentative assignments for the synthesized ligand H₂L and its coordination polymers were recorded as KBr disks and are shown in Table-2.

IR spectrum of ligand of TBO-2 show a broad band extended from 3300 to 2300 cm⁻¹ which might be responsible to phenolic -OH

group bonded to N atom of 8-hydroxyquinoline moieties²¹.

Several bands appeared between 1470 and 1610 cm⁻¹ region may arise from aromatic breathing and 3400 cm⁻¹ for -NH group. The IR band at 1580 cm⁻¹ (C=N of 8-quinolinol system) of TBO-2 ligand shifted to higher frequency side 1600 cm⁻¹ in the spectra of the metal complexes indicating involvement of nitrogen in the complexes formation²², whereas the band at

1450 cm^{-1} in the IR spectrum of TBQ-2 assigned to in-plane -OH deformation was shifted towards higher frequency in the spectra of the coordination polymer due to the formation of the M-O bond²³. This was further confirmed by a weak band at 1100 cm^{-1} corresponding to C-O-M stretching, while bands around 710 and 550 cm^{-1} correspond to the N \rightarrow M vibration²⁴.

¹H NMR Analysis

The structure of the ligand (TBQ-2) was characterized by ¹H NMR spectrum in DMSO- d₆ system.

NMR

(DMSO) 6.8 – 8.9 ppm (15H)- Multiplet Aromatic

5.3 ppm (1H)- Singlet (OH)

4.0 ppm (1H) - Singlet (NH)

Magnetic Measurements

Magnetic moments of coordination polymers are given in Table-2. The diffuse electronic spectrum of Cu⁺² complex shows two broad bands, 15590 and 22500 cm^{-1} . The first band may be due to a ²E_g \rightarrow ²T_{2g} transition, while the second band may be due to charge transfer. The first band shows structures suggesting a distorted octahedral structure for the Cu⁺² metal complex^{25,26}. The Co⁺² metal complex gives rise to two absorption bands at 21270, 18600 and 11950 cm^{-1} , which can be assigned ⁴T_{1g}(F) \rightarrow ⁴A_{2g}, ⁴T_{1g}(F) \rightarrow ⁴T_{1g}(P) and ⁴T_{1g}(F) \rightarrow ⁴T_{2g} transitions, respectively. These absorption bands and the μ_{eff} value indicate octahedral configuration of the Co⁺² metal complex^{27,28}. The spectrum of Mn⁺² polymeric complex comprised three bands at 14800 cm^{-1} , 17620 cm^{-1} and 22450 cm^{-1} . These bands may be assigned to ⁶A_{1g} \rightarrow ⁶A_{1g}(⁴E_g), ⁶A_{1g} \rightarrow ⁴T_{2g}(⁴G) and ⁶A_{1g} \rightarrow ⁴T_{1g}(⁴G) transitions, respectively. The high intensity of the bands also suggests that they may have some charge transfer character. The magnetic moment is found to be lower than normal range. In the absence of low temperature measurement of magnetic moment, it is difficult to attach any significance to this. As the spectrum of the metal

complex of Ni⁺² show three distinct bands at 22580 cm^{-1} , 15100 cm^{-1} and 8300 cm^{-1} are assigned as ³A_{2g} \rightarrow ³T_{1g}(P), ³A_{2g} \rightarrow ³T_{1g}(F) and ³A_{2g} \rightarrow ³T_{2g} transition, respectively, suggesting the octahedral environment for Ni⁺² ion. The observed μ_{eff} values in the range 2.05–5.91 B.M are consistent with the above moiety^{29,30}.

Thermal Studies

The TGA data for the Co-ordination polymers samples at different temperatures indicate that the degradation of the co-ordination polymers is noticeable beyond 310^o C. The rate of degradation becomes a maximum at a temperature between 400 and 500^o C. This may be due to acceleration by metal oxides, which form in situ. Each polymer lost about 60% of its weight when heated up to 690^o C. Inspection of the thermograms of all coordinated polymer samples revealed that all samples suffered appreciable weight loss in the range of 150 to 280^o C. This may be due to the presence of a coordinated water molecule.

Antibacterial(Antimicrobial) Activities

The increase in antimicrobial activity may be considered in light of Overtone's concept^{31,32} and Tweedy's chelation theory^{33,34}. On complication, the polarity of the metal ion will be reduced to a greater extent due to the overlap of the ligand orbital and partial sharing of the positive charge of the metal ion with donor groups. Further, it increases the delocalization of -electrons over the whole chelate ring and enhances the lipophilicity of the coordination polymers. This increased lipophilicity enhances the penetration of the coordination polymer into lipid membranes and blocks the metal binding sites in the enzymes of microorganisms. These coordination polymers also disturb the respiration process of the cell and thus block the synthesis of proteins, which restricts further growth of the organisms. The antibacterial and antifungal data obtained from analysis are shown in Table-3 and Table-4.

Table 3: Antibacterial activity of coordination Polymers

Compounds	Gram +ve		Gram -ve		
	<i>Bacillus subtilis</i>	<i>Staphylococcus Aureus</i>	<i>klebsiella promioe</i>	<i>Salmonella typhi</i>	<i>E.coli</i>
TBQ-2	33	31	29	30	30
(Cu TBQ-2 (H ₂ O) ₂) _n	40	44	40	40	37
(Co TBQ-2 (H ₂ O) ₂) _n	42	40	35	34	35
(Ni TBQ-2 (H ₂ O) ₂) _n	43	38	37	36	37
(Mn TBQ-2 (H ₂ O) ₂) _n	44	42	36	38	30
(Zn TBQ-2 (H ₂ O) ₂) _n	40	42	38	43	35

Table 4: Antifungal activity of coordination Polymers

Compounds	Zone of Inhibition at 1000 ppm (%)				
	<i>Aspergillus Niger</i>	<i>Candida albicans</i>	<i>Trichoderma harsianum</i>	<i>Mucor mucedo</i>	<i>Botrytis cinerea</i>
TBQ-2	24	27	25	24	27
(Cu TBQ-2 (H ₂ O) ₂) _n	32	42	41	35	25
(Co TBQ-2 (H ₂ O) ₂) _n	30	29	30	28	30
(Ni TBQ-2 (H ₂ O) ₂) _n	30	29	30	27	27
(Mn TBQ-2 (H ₂ O) ₂) _n	30	28	32	28	32
(Zn TBQ-2 (H ₂ O) ₂) _n	30	25	30	25	28

Coordination polymers exhibit higher biocidal activity as compared with the free ligands; from the comparative analysis shown in Table 3 and Table 4, respectively, it is observed that all the coordination polymer are more potent biocidals than the free ligands. From the data obtained it is clear that Cu (II) is highly active among the coordination polymer of the respective metal.

CONCLUSION

The results at present work show the following conclusions. The design synthesis of new bis-5-amino salicylic acid have been successfully demonstrated. FT-IR, ¹H-NMR spectral studies. We have synthesized a series of coordination polymer from these bis-ligands with transition metals have been prepared and characterized for their spectral and magnetic properties. All the synthesized coordination polymer compounds were monitored for their antimicrobial activity. The coordination polymers are toxic for gram-negative bacteria (*E.coli*, *salmonella typhi* and *klebsiella promioe*) and gram-positive bacteria (*Bacillus subtilis* and *staphylococcus aureus*), and plant pathogenic organisms (fungi) used were *Aspergillus niger*, *Candida albicans*, *Trichoderma harsianum*, *Mucor mucedo*, and *Botrytis cinerea* microorganisms. It is found that the coordination polymer were more toxic for one or more bacterial strains, thus introducing a novel class of metal-based bactericidal agents. The information as octahedral geometry of the coordination polymer was obtained from their electronic and magnetic moment values.

ACKNOWLEDGEMENT

We are grateful to the Principal, Shri R.K.Parikh Arts and Science College Petlad for providing the necessary research facilities.

REFERENCES

- Smolin EM, Rapopret L and S-Triazine and derivatives interscience: New York; 1954.
- Halverson F, Hirt R. J Chem Phys. 1951;19:711.
- Kaliyappan T and Kannan P. Prog Polym Sci. 2000;25(3):343.
- Vogel's AI: A Textbook of Quantitative Chemical Analysis, Revised by J. Besselt, R. C. Denny, J.H. Jeffery and J. Mendham, ELBS Ed., 1996, 5.
- Ivanor VM and Metkina TF. Ah Anal Khim. 1978;33:2426.
- Burckhalter JH, Stephars VC, Searberough HC, Briniger WS and Edergton WE. J Am Chem Soc. 1954;76: 4902.
- Vogel C, Heinz W and Brazil Pat. 78,05,009, 1977; Ger Pat. 1977;2:832,758,.
- Burckhalter JH, Leib R and Eswaran. JOrg Chem. 1961;26:4078.
- Horowitz H and Perros JP. J Inorg Nucl Chem. 1964;26:139.
- Bailer CJ, Judd JR and McLean ML. Coordination Polymers (WADC Technical Reports). 1959;116:58-51 lpar;Part II).
- Patel RD, Patel SR and Patel HS. Eur Polym J. 1987;23:229.
- Shah TB, Patel HS, Dixit RB and Dixit BC. Int J Polym Anal and Charact. 2003;8:369.
- Chaudhari Ankita I and Chaudhari JA. Der Pharma Chemica. 2013;5(1):150-155.
- Xian Ren C, Yuqi F, Hisanori I, Kazuhisa H and Kousaburo O. Analytical Sci. 1995; 11:313. Bis Bidentate Ligand 527 Downloaded By: [INFLIBNET India Order] At: 03:55 18 June 2011.
- Abraham W, Abraham D, Guy R and Abraham P. Reactive Polymers, Ion Exchangers, Sorbents. 1984;2:301.
- Patel KD, Pachani SC and Dixit RB. Int J Inorganic and Orgeno Metallic Polymers 2003.
- Patel HS and Patel VK. Indian J Hetrocycl Chem. 2003;12:253.
- Vogel AI. Textbook of Quantitative Chemical Analysis, 4th ed.; ELBS:London,1978.
- Murrey PR, Baran EJ, Pfuller MA, Tenovov FC and Yolken RH. An Antimicrobial Agent and Susceptibility

- Testing; American Soc Microbiology. Washington, DC, 1995;1327.
20. Phillips JP, Elbinger RL and Merritt LL. *J Am Chem Soc.* 1949;71:3984.
 21. Bellamy LJ. *Infrared Spectra of Coordination polymer Molecules*, Chapman and Hall, London, 1957.
 22. Parekh HM, Panchal PK and Patel MN. *J Therm Anal Cal.* 2006;86:803.
 23. Masoud MS, Amira MF, Ramadan AM and El-Ashry GM. *Spectrochim Acta Part A.* 2008;69:30.
 24. Satpathy KC, Pande AK and Mishra RP. *Synth React Inorg Met Org Chem.* 1991;21:531.
 25. Cotton FA and Wilkinson G. *Advanced Inorganic Chemistry*, 5th Ed., Wiley, New York, 1988.
 26. Pancholi HB, Patel MM. *J Polym Mater.* 1996;13:261–267.
 27. Singh DP, Shishodia N, Yadav BP and Rana VB. *J Indian Chem Soc.* 2004;81: 287-290.
 28. Bull AS, Martin RB and William RJP. in "Electronic Aspects of Biochemistry." ed. B. Pullmann, Academic, New York 1964.
 29. Figgis BN and Lewis J. *The Magneto Chemistry of Coordination polymer in Modern Coordination Chemistry*, Interscience, New York, 1960.
 30. Williams JO. *Adv Phys Org Chem.* 1979; 15:159.
 31. Patel IJ and Vohra IM. *E Journal of Chemistry.* 2006;3(2):110-116.
 32. Harsfall JG. *Quantitative bioassay of fungicides in the laboratory.* *Bot Rev* 1945;11:357-397.
 33. Parekh HM, Pansuriya PB and Patel MN. *Characterization and antifungal study of genuine oxovanadium(IV) mixed-ligand complexes with Schiff bases.* *Polish J.Chem.* 2005;79:1843-1851.
 34. Kharadi GJ and Patel KD. *Antibacterial, spectral and thermal aspects of drug based- Cu(II) mixed ligand complexes.* *Appl Organomet Chem.* 2009;391.